


Library
National Wetlands Research Center
U. S. Fish and Wildlife Service
700 Cajundome Boulevard
Lafayette, La. 70506

Biological Services Program

FWS/OBS-82/10.8
FEBRUARY 1982

HABITAT SUITABILITY INDEX MODELS: BLUEGILL



U. S. Fish and Wildlife Service

U. S. Department of the Interior

SK
361
.U54
no. 82-
10.8

The Biological Services Program was established within the U.S. Fish and Wildlife Service to supply scientific information and methodologies on key environmental issues that impact fish and wildlife resources and their supporting ecosystems. The mission of the program is as follows:

- To strengthen the Fish and Wildlife Service in its role as a primary source of information on national fish and wildlife resources, particularly in respect to environmental impact assessment.
- To gather, analyze, and present information that will aid decisionmakers in the identification and resolution of problems associated with major changes in land and water use.
- To provide better ecological information and evaluation for Department of the Interior development programs, such as those relating to energy development.

Information developed by the Biological Services Program is intended for use in the planning and decisionmaking process to prevent or minimize the impact of development on fish and wildlife. Research activities and technical assistance services are based on an analysis of the issues, a determination of the decisionmakers involved and their information needs, and an evaluation of the state of the art to identify information gaps and to determine priorities. This is a strategy that will ensure that the products produced and disseminated are timely and useful.

Projects have been initiated in the following areas: coal extraction and conversion; power plants; geothermal, mineral and oil shale development; water resource analysis, including stream alterations and western water allocation; coastal ecosystems and Outer Continental Shelf development; and systems inventory, including National Wetland Inventory, habitat classification and analysis, and information transfer.

The Biological Services Program consists of the Office of Biological Services in Washington, D.C., which is responsible for overall planning and management; National Teams, which provide the Program's central scientific and technical expertise and arrange for contracting biological services studies with states, universities, consulting firms, and others; Regional Staffs, who provide a link to problems at the operating level; and staffs at certain Fish and Wildlife Service research facilities, who conduct in-house research studies.

HABITAT SUITABILITY INDEX MODELS: BLUEGILL

by

Robert J. Stuber
Habitat Evaluation Procedures Group
Western Energy and Land Use Team
U.S. Fish and Wildlife Service
Drake Creekside Building One
2625 Redwing Road
Fort Collins, Colorado 80526

Glen Gebhart
and
O. Eugene Maughan
Oklahoma Cooperative Fishery Research Unit
Oklahoma State University
Stillwater, Oklahoma 74074

Western Energy and Land Use Team
Office of Biological Services
Fish and Wildlife Service
U.S. Department of the Interior
Washington, D.C. 20240

This report should be cited as:

Stuber, R. J., G. Gebhart, and O. E. Maughan. 1982. Habitat suitability index models: Bluegill. U.S.D.I. Fish and Wildlife Service. FWS/OBS-82/10.8. 26 pp.

PREFACE

The habitat use information and Habitat Suitability Index (HSI) models presented in this document are an aid for impact assessment and habitat management activities. Literature concerning a species' habitat requirements and preferences is reviewed and then synthesized into HSI models, which are scaled to produce an index between 0 (unsuitable habitat) and 1 (optimal habitat). Assumptions used to transform habitat use information into these mathematical models are noted, and guidelines for model application are described. Any models found in the literature which may also be used to calculate an HSI are cited, and simplified HSI models, based on what the authors believe to be the most important habitat characteristics for this species, are presented.

Use of the models presented in this publication for impact assessment requires the setting of clear study objectives and may require modification of the models to meet those objectives. Methods for reducing model complexity and recommended measurement techniques for model variables are presented in Appendix A.

The HSI models presented herein are complex hypotheses of species-habitat relationships, not statements of proven cause and effect relationships. Results of model performance tests, when available, are referenced; however, models that have demonstrated reliability in specific situations may prove unreliable in others. For this reason, the FWS encourages model users to convey comments and suggestions that may help us increase the utility and effectiveness of this habitat-based approach to fish and wildlife planning. Please send comments to:

Habitat Evaluation Procedures Group
Western Energy and Land Use Team
U.S. Fish and Wildlife Service
2625 Redwing Road
Ft. Collins, CO 80526

CONTENTS

	<u>Page</u>
PREFACE	iii
ACKNOWLEDGEMENTS	vi
HABITAT USE INFORMATION	1
General	1
Age, Growth, and Food	1
Reproduction	1
Specific Habitat Requirements	1
HABITAT SUITABILITY INDEX (HSI) MODELS	3
Model Applicability	3
Model Description - Riverine	3
Model Description - Lacustrine	6
Suitability Index (SI) Graphs for Model Variables	7
Riverine Model	14
Lacustrine Model	15
Interpreting Model Outputs	16
ADDITIONAL HABITAT MODELS	16
Model 1	16
Model 2	23
Model 3	23
REFERENCES CITED	23

ACKNOWLEDGEMENTS

We would like to thank Armond Lemke, Richard Anderson, William Reynolds, and Martha Casterlin for reviewing the manuscript. Their review contributions are gratefully acknowledged, but the authors accept full responsibility for the contents of the document. Word processing was provided by Dora Ibarra and Carolyn Gulzow. The cover of this document was illustrated by Jennifer Shoemaker.

BLUEGILL (Lepomis macrochirus)

HABITAT USE INFORMATION

General

The bluegill (Lepomis macrochirus) is native from the Lake Champlain and southern Ontario region through the Great Lakes to Minnesota, and south to northeastern Mexico, the Gulf States, and the Carolinas (Scott and Crossman 1973). The species has been widely introduced outside its native range (Pflieger 1975). Three subspecies are currently recognized: L. m. macrochirus (northcentral United States), L. m. speciosus (Texas and northern Mexico), and L. m. purpurascens (Atlantic and Gulf States) (Hubbs and Lagler 1958).

Age, Growth, and Food

Bluegills usually mature at age I or II (Schloemer 1939; James 1946; Cross 1951). The maximum known age is 11 years, but most live 1 to 4 years (Schloemer 1939; Carlander 1977). Maximum recorded size is 39 cm and 2.1 kg (Emig 1966).

Bluegills are opportunistic feeders which can alter their diet according to food availability (Keast and Webb 1966). Fry feed primarily on zooplankton and small insects (Werner 1969). Juveniles and adults feed on zooplankton, aquatic and terrestrial insects, and some plant materials (Scidmore and Woods 1960; Emig 1966; Scott and Crossman 1973).

Reproduction

Bluegills are repeat spawners and the spawning season may extend from spring through summer (Anderson, pers. comm.). Spawning occurs from 17 to 31° C, with peak spawning at 24-27° C (Clugston 1966; Emig 1966; Scott and Crossman 1973; Kitchell et al. 1974; Pflieger 1975). Bluegills are guarding, nest building lithophils (Balon 1975). Nests are usually found in quiet, shallow (1-3 m) water (Swingle and Smith 1943). Although spawning will occur over almost any substrate, fine gravel or sand is preferred (Stevenson et al. 1969; Pflieger 1975). Incubation time ranges from 1.5 to 5 days, depending on ambient water temperature (Morgan 1951; Childers 1967; Heckman 1969; Hall et al. 1970; Merriner 1971).

Specific Habitat Requirements

Bluegills are most abundant along shoreline areas in lentic and lentic-type environments such as ponds, lakes, reservoirs, and large low velocity streams (Whitmore et al. 1960). In riverine habitats, bluegills are mostly restricted to areas of low velocity (Hubbs and Lagler 1958). Adult bluegills were captured primarily in backwater areas of the Missouri River (Kallemyn and Novotny 1977). Hardin and Bovee (1978) developed probability of use curves showing that adults prefer current velocities < 10 cm/sec but will tolerate up to 45 cm/sec. Abundance has been positively correlated to a high percent (> 60%) pool area and negatively correlated to a high percent riffle/run area

(Moyle and Nichols 1973). Optimal stream gradient (≤ 0.5 m/km) is based on the preference for low gradient, lentic-type waters (Trautman 1957).

Optimal lacustrine habitat is characterized by fertile lakes, ponds, and reservoirs with extensive ($\geq 20\%$ of lacustrine surface area) littoral areas (Emig 1966; Scott and Crossman 1973). However, deeper areas are also required for overwintering and retreat from the summer heat (Scott and Crossman 1973). Jenkins (1976) reported a significant positive correlation between TDS levels of 100-350 ppm and sportfish (including sunfishes) standing crops in a group of predominately southeastern reservoirs.

Cover in both lacustrine and riverine habitats in the form of submerged vegetation or logs and brush is utilized by the species, especially juveniles and small adults (Moyle and Nichols 1973; Scott and Crossman 1973). However, an excessive abundance of vegetation can inhibit utilization of prey by bluegills. Populations of stunted individuals have been associated with an excessive amount of aquatic vegetation which may inhibit the utilization of bluegills as prey (Anderson, pers. comm.). Bluegills also nest in unvegetated areas (Weaver and Ziebell 1976). Lack of cover may also be a problem (Anderson, pers. comm.).

Water quality criteria for bluegills in both riverine and lacustrine habitats are outlined as follows: optimal growth and reproductive potential occurs in waters of low to moderate turbidities (< 50 ppm) (Buck 1956; Hastings and Cross 1962; Shireman 1968). Bluegills can tolerate a pH range of 4.0 to 10.3 (Trama 1954; Ultsch 1978) but pH levels at these extremes have caused at last partial kills (Calabrese 1969). Optimal levels are 6.5-8.5 based on Stroud's (1967) criteria for freshwater fish. Bluegills can tolerate dissolved oxygen levels < 1.0 mg/l for short durations (Baker 1941; Cooper and Washburn 1946; Moss and Scott 1961; Petrosky and Magnuson 1973) but will avoid levels of 1.5-3.0 mg/l (Whitmore et al. 1960). Optimal levels are > 5.0 mg/l (Petit 1973). Bluegills will not tolerate salinities > 5.6 ppt (Kilby 1955), and prefer salinity levels < 3.6 ppt (Tebo and McCoy 1964; Carver 1967).

Adult. Optimal growth of adult bluegills occurs near 27° C (Anderson 1959). No growth occurs below 10° C or above 30° C (Anderson 1959; Emig 1966). The reported ultimate upper incipient lethal temperature for bluegill is 35° C (Reynolds and Casterlin 1976).

Embryo. Optimal temperatures for successful embryo development are $22-27^{\circ}$ C, and development will occur from $22-34^{\circ}$ C (Banner and Van Arman 1973). Optimal current velocities are < 7.5 cm/sec, and embryos are not found at current velocities > 30 cm/sec (Hardin and Bovee 1978). Because bluegill spawn at 1-3 m depth (Swingle and Smith 1943), reservoir drawdown during spawning should not exceed 3 m during spring and summer.

Fry. Optimal temperatures for fry are $25-32^{\circ}$ C (Hardin and Bovee 1978). Fry will not survive temperatures below 11° C or above 34° C (Banner and Van Arman 1973). Optimal current velocities are < 5 cm/sec; fry are not found in areas with velocities greater than about 7.5 cm/sec (Kallemyn and Novotny 1977; Hardin and Bovee 1978).

Juvenile. The highest specific growth rate of juvenile bluegill occur in waters of 30° C and the growth range is $22-34^{\circ}$ C (Lemke 1977). Preferred

current velocities are < 5 cm/sec; juveniles are not found in areas with velocities greater than about 15 cm/sec (Kallemyn and Novotny 1977; Hardin and Bovee 1978).

HABITAT SUITABILITY INDEX (HSI) MODELS

Model Applicability

Geographic area. The model is applicable wherever bluegills occur in North America. The standard of comparison for each individual variable suitability index is the optimum value of the variable that occurs anywhere within this region. Therefore, the model will never provide an HSI of 1.0 when applied to water bodies in the North where temperature related variables do not reach the optimum values found in the South.

Season. The model provides a rating for a riverine or lacustrine habitat based on its ability to support all life stages of bluegills through all seasons of the year.

Cover types. The model is applicable in riverine and lacustrine habitats as described by Cowardin et al. (1979).

Minimum habitat area. Minimum habitat area is defined as the minimum area of contiguous suitable habitat that is required for a population to live and reproduce. No attempt has been made to establish a minimum habitat size for survival and growth of a bluegill population.

Verification level. The acceptance goal of the model is to produce an index between 0 and 1 which has a positive relationship to spawning success of adults and carrying capacity for fry, juveniles, and adults. In order to verify that the model output was acceptable, HSI's were calculated from sample data sets. These sample data sets and their relationship to model verification are discussed in greater detail following the presentation of the model.

Model Description - Riverine

Variables which have been shown to affect growth, survival, abundance, or other measure of well-being of bluegill are placed in the appropriate component (Figs. 1 and 2).

Food component. Percent cover (logs and other objects) (V_2) is included because this type of cover in pools is favorable prey habitat. Percent cover (vegetation) (V_3) is included as a separate variable because vegetation density can influence both feeding ability of bluegills and abundance of food. Percent pools (V_1) is included to quantify the amount of food habitat.

Cover component. Percent cover (logs and other objects) (V_2) and percent cover (vegetation) (V_3) are included because bluegills exhibit cover-seeking behavior. Percent vegetative cover is separate from other cover because too much vegetation can cause a habitat problem, while logs, brush, and other debris provide good cover.

Habitat Variables

Life Requisites




Figure 1. Tree diagram illustrating relationship of habitat variables and life requisites in the riverine model for the bluegill. Dashed line indicates optional variable in the model.

Habitat Variables

Life Requisites




Figure 2. Tree diagram illustrating relationships of habitat variables and life requisites in the lacustrine model for the bluegill. Dashed line indicates optional variable in the model.

Water quality component. Turbidity (V_6), pH (V_7), dissolved oxygen (V_8), and temperature (V_{10} , V_{12} , and V_{13}) are crucial parameters that affect development, growth, and survival. Dissolved oxygen and temperature are weighted in the model and are considered to be limiting factors. Salinity (V_9) is an optional variable since it is not considered to be a problem in most areas where bluegills are found.

Reproduction component. Temperature (V_{11}) is included because embryo survival and development depends on the temperature being warm enough for incubation and hatching. Current velocity in spawning areas (V_{15}) is included because the embryo will not survive in areas of higher velocities. Substrate composition (V_{20}) is included since bluegill show a preference for spawning over fine gravel and sand.

"Other" component. The variables in the "other" component are those which aid in describing habitat suitability for bluegills, yet are not specifically related to life requisite components already presented. Current velocity for the different life stages (V_{14} , V_{16} , and V_{17}) is included because all stages cannot tolerate swifter velocities. Stream gradient (V_{18}) is included because bluegills are most often found in lower gradient streams.

Model Description - Lacustrine

Food component. Percent cover (logs and other objects) (V_2) is included because this type of habitat promotes good habitat for foraging and food organisms. Percent cover (vegetation) (V_3) is included because, though vegetation can be a measure of lacustrine productivity, too much vegetation can seriously reduce foraging capabilities. Total dissolved solids (TDS) (V_5) is included because TDS is a measure of general lacustrine productivity and dissolved solids are a vital prerequisite for the development of the food chain. Bluegills are opportunistic feeders on whatever is abundant. pH (V_6) is included in the food component because a low pH may indicate low alkalinity and low productivity.

Cover component. Percent cover (logs and other objects) (V_2) and percent cover (vegetation) (V_3) are important since cover-seeking behavior indicates that some cover must exist for good habitat. Too much vegetative cover may indicate poor habitat. Percent littoral area (V_4) is included to quantify the amount of cover habitat.

Water quality component. See description for riverine water quality component.

Reproduction component. Temperature (V_{11}) is included because embryo survival and development is related to temperature. Reservoir drawdown (V_{19}) is included because bluegills spawn at a certain depth and eggs may be exposed if water levels drop too low (this variable is excluded in a natural lake or pond). Substrate composition (V_{20}) is included because bluegill show a preference for spawning over fine gravel and sand.


Suitability Index (SI) Graphs for Model Variables

This section contains suitability index graphs for the 20 variables described above. The "R" pertains to riverine habitat variables, and the "L" refers to lacustrine habitat variables.


Habitat Variable

Suitability Graph


R (V_1) Percent pool area during average summer flow.




R,L (V_2) Percent cover (e.g., logs, brush, and debris) within pools or littoral areas during summer.



R,L (V₃) Percent cover (aquatic vegetation, submersed, dense stands, finely divided leaves).




L (V₄) Percent littoral area during summer stratification.




L (V₅) Average TDS level during growing season.

Note: SI values should be lowered about 0.2 if ionic concentration of sulfate-chlorides exceeds that of carbonate-bicarbonates.




R,L (V₆) Maximum monthly average turbidity during average summer flow or summer stratification.



R,L (V₇) pH range during growing season.


- A) 6.5-8.5
- B) 6.0-6.5 or 8.5-9.0
- C) 5.0-6.0 or 9.0-10.0
- D) < 5.0 or > 10.0



R,L (V₈) Minimum dissolved oxygen range during summer.

- A) Seldom below 5.0 mg/l
- B) Usually between 3.0 and 5.0 mg/l
- C) Usually between 1.5 and 3.0 mg/l
- D) Often below 1.5 mg/l


Note: Lacustrine D.O. levels refer to littoral areas; riverine, pools.



R,L

(V₉)


Maximum monthly average salinity during growing season (optional).



R,L

(V₁₀)


Maximum midsummer temperature within pools or littoral areas (adult).




R,L

(V₁₁)


Average of mean weekly water temperature within pools or littoral areas during spawning (embryo).




R,L (V₁₂) Maximum early summer temperature within pools or littoral areas (fry).




R,L (V₁₃) Maximum midsummer temperature within pools or littoral areas (juvenile).




R (V₁₄) Average current velocity in pools and backwater areas during growing season (adult).




R (V₁₅) Average current velocity in spawning areas (embryo).




R (V₁₆) Average current velocity in pools and backwater areas during early summer (fry).




R (V₁₇) Average current velocity in pools and backwater areas during growing season (juvenile).



R (V₁₈) Stream gradient within representative reach.




L (V₁₉) Reservoir drawdown during spawning (embryo).



R,L (V₂₀) Substrate composition within pools or littoral areas during spawning (embryo).

- A) Fines and gravel present
- B) Fines and gravel scarce



Riverine Model

This model utilizes the life requisite approach and consists of five components: food, cover, water quality, reproduction, and other.

Food (C_F).

$$C_F = (V_1 \times V_2 \times V_3)^{1/3}$$

Cover (C_C).

$$C_C = \frac{V_2 + V_3}{2}$$

Water Quality (C_{WQ}).

$$C_{WQ} = \frac{V_6 + V_7 + 2V_8 + V_9 + 2[(V_{10} \times V_{12} \times V_{13})^{1/3}]}{7}$$

If V_8 or $(V_{10} \times V_{12} \times V_{13})^{1/3} \leq 0.4$, C_{WQ} equals the lowest of the following: V_8 , $(V_{10} \times V_{12} \times V_{13})^{1/3}$, or the above equation.

Note: V_9 may be dropped and the denominator changed to 6 if salinity is not considered to be a problem or potential problem in the study area.

Reproduction (C_R).

$$C_R = (V_{11} \times V_{15} \times V_{20})^{1/3}$$

Other (C_{OT}).

$$C_{OT} = \frac{\frac{V_{14} + V_{16} + V_{17}}{3} + V_{18}}{2}$$

HSI determination. If all component ratings > 0.4,

$$HSI = (C_F \times C_C \times C_{WQ}^2 \times C_R \times C_{OT})^{1/6}$$

If C_{WQ} or C_R are ≤ 0.4 , use lowest component rating as the species HSI.

Lacustrine Model

This model utilizes the life requisite approach and consists of four components: food, cover, water quality, and reproduction.

Food (C_F).

$$C_F = (V_2 \times V_3 \times V_5 \times V_7)^{1/4}$$

Cover (C_C).

$$C_C = (V_2 \times V_3 \times V_4^2)^{1/4}$$

Water Quality (C_{WQ}).

$$C_{WQ} = \frac{V_6 + V_7 + 2V_8 + V_9 + 2[(V_{10} \times V_{12} \times V_{13})^{1/3}]}{7}$$

If V_8 or $(V_{10} \times V_{12} \times V_{13})^{1/3} \leq 0.4$, C_{WQ} equals the lowest of the following: V_8 , $(V_{10} \times V_{12} \times V_{13})^{1/3}$, or the above equation.

Note: V_9 may be dropped and the denominator changed to 6 if salinity is not considered to be a problem or potential problem in the study area.

Reproduction (C_R).

$$C_R = (V_{11} \times V_{19} \times V_{20})^{1/3}$$

Note: If the lacustrine environment is a natural lake or pond, V_{19} will not be applicable. Thus,

$$C_R = (V_{11} \times V_{20})^{1/2} \text{ in a natural lake or pond}$$

HSI determination. If all component ratings > 0.4,

$$HSI = (C_F \times C_C \times C_{WQ}^2 \times C_R)^{1/5}$$

If C_{WQ} or $C_R \leq 0.4$, use lowest component ratings as the species HSI.

Sources of data and assumptions made in developing the suitability indices are presented in Table 1.

Sample data sets for the above riverine and lacustrine HSI models are listed in Tables 2 and 3. The data sets are not actual field measurements but represent combinations that could occur in a riverine or lacustrine habitat. The HSI's calculated from the data reflect what the carrying capacity trends would be in riverine and lacustrine habitats with the listed characteristics. Thus, the model meets the acceptance goal of producing an index between 0 and 1 which is believed to have a positive relationship to the spawning success of adults and carrying capacity of fry, juvenile, and adult bluegill.

Interpreting Model Outputs

Habitats with an HSI of 0 may contain some bluegills; habitats with a high HSI may contain few. The bluegill HSI determined by use of these models will not necessarily represent the population of bluegill in the study area. This is because the standing crop does not totally depend on the ability of the habitat to meet all life requisite requirements of the species. If the model is a good representation of bluegill riverine or lacustrine habitat, the model should be positively correlated with long term average population levels in areas where population levels are determined primarily by habitat related factors. However, this has not been tested. The proper interpretation of the HSI produced by the model is one of comparison. If two habitats have different HSI's, the one with the higher HSI should have the potential to support more fish than the one with the lower HSI, given the model assumptions have not been violated.

ADDITIONAL HABITAT MODELS

Model 1

Optimal riverine habitat for bluegills is characterized by the following conditions, assuming water quality is adequate: large, low gradient (< 0.5 m/km) streams; warm water temperatures (> 20° C); sluggish current velocities (< 5 cm/sec); clear water (< 50 ppm suspended solids); and an abundance of bottom cover within pool areas.

$$HSI = \frac{\text{number of above criteria present}}{5}$$

Table 1. Data sources for bluegill suitability indices.

Variable and source	Assumption
V ₁ Moyle and Nichols 1973	The amount of pool area that is correlated to a high abundance of bluegills is optimum.
V ₂ Moyle and Nichols 1973 Scott and Crossman 1973 Pflieger 1975	The percent cover (logs and other objects) where bluegill are most abundant is optimum.
V ₃ Moyle and Nichols 1973 Scott and Crossman 1973 Weaver and Ziebell 1976 Anderson, pers. comm.	The percent cover (vegetation) that is associated with abundant fish is optimum. Not enough vegetative cover or too much vegetative cover are suboptimum, since the former restricts the food availability and the latter restricts foraging capabilities.
V ₄ Emig 1966 Scott and Crossman 1973	Since the bluegill inhabits shallow vegetated areas, a certain percentage of littoral area must exist for habitat to be suitable. Since bluegills require deeper water in winter and to get away from summer heat, too much littoral area would be suboptimum to unsuitable.
V ₅ Jenkins 1976	TDS levels associated with high standing crops are optimum. Levels that reduce food availability are suboptimum to unsuitable.
V ₆ Buck 1956 Trautman 1957 Hastings and Cross 1962 Shireman 1968 Pflieger 1975	Turbidity levels where growth rates are fastest are optimum. Levels that retard growth and development and that adversely affect reproduction are suboptimum to unsuitable.
V ₇ Trama 1954 Stroud 1967 Calabrese 1969 Ultsch 1978	pH levels that promote good production and maximum survival are optimum. Levels that reduce reproductive capabilities and feeding are suboptimum to unsuitable.
V ₈ Cooper and Washburn 1946 Whitmore et al. 1960 Doudoroff and Shumway 1970 Petit 1973	D.O. levels where survival is maximum and development is normal are optimum. Levels causing stress reactions are suboptimum. Levels that are tolerated for short durations are unsuitable.

Table 1. (continued)

Variable and source	Assumption
V ₉ Kilby 1955 Tebo and McCoy 1964 Carver 1967	Salinity levels that promote successful reproduction and good growth are optimum. Levels where the species does not reproduce are unsuitable.
V ₁₀ Anderson 1959 Emig 1966	Temperatures that promote maximum growth are optimum. Temperatures where no growth occurs are unsuitable.
V ₁₁ Clugston 1966 Emig 1966 Banner and Van Arman 1973 Scott and Crossman 1973 Kitchell et al. 1974 Pflieger 1975	Temperatures where embryo development is successful and normal and survival is maximum are optimum. Temperatures where survival is reduced but development may occur are suboptimum. Temperatures where no development occurs are unsuitable.
V ₁₂ Banner and Van Arman 1973 Hardin and Bovee 1978	Temperatures that reach levels where maximum growth occurs are optimum. Temperatures where the species does not survive are unsuitable.
V ₁₃ Lemke 1977	Temperatures that reach levels where maximum growth occurs are optimum. Temperatures where feeding is reduced but where growth still occurs are suboptimum. Temperatures that cause no growth or death are unsuitable.
V ₁₄ Kallemyn and Novotny 1977 Hardin and Bovee 1978	Current velocities where bluegills are most often collected are optimum. Velocities where the species is seldom or never found are suboptimum to unsuitable.
V ₁₅ Same as V ₁₄	Same as V ₁₄
V ₁₆ Same as V ₁₄	Same as V ₁₄
V ₁₇ Same as V ₁₄	Same as V ₁₄
V ₁₈ Trautman 1957	Stream gradients where bluegills are collected in abundant numbers are optimum. Gradients where the fish occur in fewer numbers or are absent are suboptimum to unsuitable.

Table 1. (concluded)

Variable and source	Assumption
V ₁₉ Swingle and Smith 1943	Because bluegill spawn at specific depths, stable water levels are optimum. Any reservoir drawdown would be suboptimum to unsuitable.
V ₂₀ Stevenson et al. 1969 Pflieger 1975	Substrates that the species prefers are optimum. Almost any other substrate has high suitability.

Table 2. Sample data sets using riverine HSI model.

Variable		Data set 1		Data set 2		Data set 3	
		Data	SI	Data	SI	Data	SI
% pools	V ₁	20	0.3	35	0.6	70	1.0
% cover (logs, brush)	V ₂	5	0.4	10	0.6	20	1.0
% cover (vegetation)	V ₃	5	0.3	10	0.6	25	1.0
Turbidity	V ₆	140	0.5	70	0.9	110	0.7
pH	V ₇	Class A	1.0	Class A	1.0	Class A	1.0
Dissolved oxygen (mg/l)	V ₈	Class C	0.4	Class A	1.0	Class B	0.7
Salinity (ppt)	V ₉	1.2	1.0	2.0	1.0	0.8	1.0
Temperature - adult (°C)	V ₁₀	24	0.9	26	1.0	27	1.0
Temperature - embryo (°C)	V ₁₁	18.5	0.1	22	1.0	24	1.0
Temperature - fry (°C)	V ₁₂	22	0.8	24	0.9	27	1.0
Temperature - juvenile (°C)	V ₁₃	22	0.5	24	0.6	27	0.8
Current velocity - adult (cm/sec)	V ₁₄	20	0.7	30	0.4	9	1.0
Current velocity - embryo (cm/sec)	V ₁₅	28	0.3	30	0.2	14	0.8
Current velocity - fry (cm/sec)	V ₁₆	20	0.0	30	0.0	10	0.1
Current velocity - juvenile (cm/sec)	V ₁₇	20	0.0	30	0.0	9	0.6
Stream gradient (m/km)	V ₁₈	2	0.5	2.7	0.3	0.7	0.9
Substrate comp.	V ₂₀	Class B	0.7	Class A	1.0	Class A	1.0

Table 2. (concluded)

Variable	Data set 1		Data set 2		Data set 3	
	Data	SI	Data	SI	Data	SI
<u>Component SI</u>						
$C_F =$		0.33		0.60		1.00
$C_C =$		0.35		0.60		1.00
$C_{WQ} =$		0.67		0.93		0.85
$C_R =$		0.28		0.58		0.93
$C_{OT} =$		0.37		0.22		0.73
HSI =		0.28		0.58		0.89

Table 3. Sample data sets using lacustrine HSI model.

Variable		Data set 1		Data set 2		Data set 3	
		Data	SI	Data	SI	Data	SI
% cover (logs, brush)	V ₂	100	0.2	10	0.6	50	1.0
% cover (vegetation)	V ₃	5	0.4	35	0.9	25	1.0
% littoral area	V ₄	8	0.4	17	0.8	60	1.0
TDS (ppm)	V ₅	50	0.4	10	0.1	200	1.0
Turbidity (ppm)	V ₆	15	1.0	10	1.0	90	0.8
pH	V ₇	Class A	1.0	Class A	1.0	Class A	1.0
Dissolved oxygen (mg/l)	V ₈	Class A	1.0	Class A	1.0	Class A	1.0
Salinity (ppt)	V ₉	0.4	1.0	0.2	1.0	0.5	1.0
Temperature - adult (°C)	V ₁₀	24	0.9	24	0.9	21	0.7
Temperature - embryo (°C)	V ₁₁	19	0.3	21	0.7	24	1.0
Temperature - fry (°C)	V ₁₂	21	0.7	22	0.8	28	1.0
Temperature - juvenile (°C)	V ₁₃	20	0.4	22	0.5	28	0.9
Reservoir drawdown (m)	V ₁₉	4	0.1	-	-	-	-
Substrate	V ₂₀	Class B	0.7	Class A	1.0	Class A	1.0
<u>Component SI</u>							
C _F =			0.42		0.48		1.00
C _C =			0.34		0.77		1.00
C _{WQ} =			0.89		0.87		0.93
C _R =			0.28		0.84		1.00
HSI =			0.28		0.77		0.97

Model 2

Optimal lacustrine habitat for bluegill sunfish is characterized by the following conditions, assuming water quality is adequate: fertile lakes, reservoirs, and ponds (TDS levels 100-350 ppm); extensive littoral areas ($\geq 20\%$ surface area); maximum water temperature $> 20^{\circ}\text{C}$; and clear water (< 50 ppm suspended solids).

$$\text{HSI} = \frac{\text{number of above criteria present}}{4}$$

Model 3

Use the regression models for bluegill standing crop in reservoirs presented by Aggus and Morais (1979) to calculate an HSI.

REFERENCES CITED


- Aggus, L. R., and D. I. Morais. 1979. Habitat suitability index equations for reservoirs based on standing crop of fish. Natl. Reservoir Res. Proj., Rep. to U.S. Fish Wildl. Serv., Hab. Eval. Program, Ft. Collins, Colorado. 120 pp.
- Anderson, R. O. 1959. The influence of season and temperature on growth of the bluegill, Lepomis macrochirus. Ph.D. Thesis, Univ. Michigan, Ann Arbor. 133 pp.
- _____. 1981. Personal communication. Missouri Cooperative Fishery Unit, Columbia, Missouri.
- Balon, E. K. 1975. Reproductive guilds of fishes: a proposal and definition. J. Fish. Res. Board Can. 32:821-864.
- Baker, C. L. 1941. Effects of fish gulping atmospheric air from waters of various carbon dioxide tensions. J. Tenn. Acad. Sci. 17:39-50.
- Banner, A., and J. A. Van Arman. 1973. Thermal effects on eggs, larvae and juvenile bluegill sunfish. U.S. Environ. Protection Agency, Ecol. Res. Ser. EPA-R3-73-041. 111 pp.
- Buck, D. H. 1956. Effects of turbidity on fish and fishing. Trans. N. Am. Wildl. Conf. 21:249-261.
- Calabrese, A. 1969. Effect of acids and alkalis on survival of bluegills and largemouth bass. U.S. Bur. Sport Fish Wildl. Tech. Pap. 42. 10 pp.
- Carlander, K. D. 1977. Handbook of freshwater fishery biology. Vol. II. Iowa State Univ. Press, Ames. 431 pp.

- Carver, D. C. 1967. Distribution and abundance of the centrarchids in the recent delta of the Mississippi River. Proc. Annu. Conf. Southeast. Assoc. Game Fish Commissioners 20:390-404.
- Childers, W. F. 1967. Hybridization of four species of sunfishes (Centrarchidae). Ill. Nat. Hist. Surv. Bull. 29:159-214.
- Clugston, J. P. 1966. Centrarchid spawning in the Florida everglades. J. Fla. Acad. Sci. 29:137-144.
- Cooper, G. P., and G. N. Washburn. 1946. Relation of dissolved oxygen to winter mortality of fish in Michigan lakes. Trans. Am. Fish. Soc. 76:23-33.
- Cross, F. B. 1951. Early limnological and fish population conditions of Canton Reservoir, Oklahoma with species reference to carp, channel catfish, largemouth bass, green sunfish and bluegill, and fishery management recommendations. Ph.D. Thesis, Oklahoma Agric. Mech. Coll., Stillwater. 92 pp.
- Cowardin, L. M., V. Carter, F. C. Golet, and E. J. LaRoe. 1979. Classification of wetlands and deepwater habitats of the United States. USDI Fish and Wildl. Serv., FWS/OBS-79/31. 103 pp.
- Doudoroff, P., and D. L. Shumway. 1970. Dissolved oxygen requirements of freshwater fishes. Food and Agric. Organ. Fish Tech. Pap. 86. 291 pp.
- Emig, J. W. 1966. Bluegill sunfish. Pages 375-392 in A. Calhoun, ed. Inland fisheries management. Calif. Dept. Fish Game.
- Hall, D. J., W. E. Cooper, and E. E. Werner. 1970. An experimental approach to the production dynamics and structure of freshwater animal communities. Limnol. Oceanog. 15(6):839-928.
- Hardin, T., and K. Bovee. 1978. Bluegill sunfish. Unpublished Data, Instream Flow Group, Western Energy and Land Use Team. U.S. Fish Wildl. Serv., Ft. Collins, Colorado.
- Hastings, C. E., and F. B. Cross. 1962. Farm ponds in Douglas County, Kansas. Univ. Kansas Mus. Nat. Hist. Misc. Publ. 29. 21 pp.
- Heckman, J. R. 1969. Embryological comparison of Lepomis macrochirus x macrochirus and Lepomis macrochirus x gibbosus. Trans. Am. Fish. Soc. 98:669-675.
- Hubbs, C. L., and K. F. Lagler. 1958. Fishes of the Great Lakes region. Univ. Michigan Press, Ann Arbor. 213 pp.
- James, M. F. 1946. Histology of gonadal changes in the bluegill, Lepomis macrochirus Rafinesque and the largemouth bass, Huro salmoides Lacepede. J. Morph. 79:63-92.

- Jenkins, R. M. 1976. Prediction of fish production in Oklahoma reservoirs on the basis of environmental variables. *Ann. Okla. Sc.* 5:11-20.
- Kallemyn, L. W., and J. F. Novotny. 1977. Fish and fish food organisms in various habitats of the Missouri River in South Dakota, Nebraska, and Iowa. U.S.D.I. Fish Wildl. Serv. FWS/OBS-77/25.
- Keast, A., and D. Webb. 1966. Mouth and body form relative to feeding ecology in the fish fauna of a small lake, Lake Opinicon, Ontario. *J. Fish. Res. Board Can.* 23:1845-1867.
- Kilby, J. D. 1955. The fishes of two gulf coast marsh areas of Florida. *Tulane Stud. Zool.* 2:175-247.
- Kitchell, J. F., J. F. Koonce, R. V. O'Neill, H. H. Shugart, J. J. Magnuson, and R. S. Booth. 1974. Model of fish biomass dynamics. *Trans. Am. Fish. Soc.* 103:786-798.
- Lemke, A. E. 1977. Optimum temperature for growth of juvenile bluegills. *Prog. Fish-Cult.* 39:55-57.
- Merriner, J. V. 1971. Development of intergeneric centrarchid hybrid embryos. *Trans. Am. Fish. Soc.* 100:611-618.
- Morgan, G. D. 1951. The life history of the bluegill sunfish, Lepomis macrochirus of Buckeye Lake (Ohio). *Dennison Univ. Bull. J. Sci. Lab.* 42(4):21-59.
- Moss, D. D., and D. C. Scott. 1961. Dissolved-oxygen requirements of three species of fish. *Trans. Am. Fish. Soc.* 90:377-393.
- Moyle, P. B., and R. D. Nichols. 1973. Ecology of some native and introduced fishes of the Sierra Nevada foothills in central California. *Copeia* 1973:478-490.
- Petit, G. D. 1973. Effects of dissolved oxygen on survival and behavior of selected fishes of western Lake Erie. *Ohio Biol. Surv. Bull.* 4:1-76.
- Petrosky, B. R., and J. M. Magnuson. 1973. Behavioral responses of northern pike, yellow perch and bluegill to oxygen concentrations under simulated winter kill conditions. *Copeia* 1973:124-133.
- Pflieger, W. L. 1975. The fishes of Missouri. Missouri Dept. Conserv., Jefferson City. 343 pp.
- Reynolds, W. W., and M. E. Casterlin. 1976. Thermal preferenda and behavioral thermoregulation in three centrarchid fishes. Pages 185-190 in G. W. Esch and R. W. McFarlane, eds. *Thermal Ecology II*. Tech. Info. Serv., Springfield, Va.
- Schloemer, C. L. 1939. The age and rate of growth of the bluegill, Helioperca macrochira (Raf.). Ph.D. Thesis, Univ. Wisconsin. 113 pp.

- Scidmore, W. J., and D. E. Woods. 1960. Some observations on competition among several species of fish for summer foods in four southern Minnesota lakes in 1955, 1956, and 1957. Minnesota Dept. Conserv., Fish and Game Invest., Fish Ser. 2:13-24.
- Scott, W. B., and E. J. Crossman. 1973. Freshwater fishes of Canada. Fish. Res. Board Can. Bull. 184. 966 pp.
- Shireman, J. V. 1968. Age and growth of bluegills, Lepomis macrochirus Rafinesque, from selected central Iowa farm ponds. Proc. Iowa Acad. Sci. 75:170-178.
- Stevenson, F., W. T. Momot, and F. J. Svoboda, III. 1969. Nesting success of the bluegill, Lepomis macrochirus Rafinesque in a small Ohio farm pond. Ohio J. Sci. 69:347-355.
- Stroud, R. H. 1967. Water quality criteria to protect aquatic life: a summary. Am. Fish. Soc. Spec. Publ. 4:33-37.
- Swingle, H. S., and E. V. Smith. 1943. Factors affecting the reproduction of bluegill bream and largemouth black bass in ponds. Alabama Polytech. Inst. Agric. Exp. Stn. Circ. 87. 8 pp.
- Tebo, L. B., Jr., and E. G. McCoy. 1964. Effects of seawater concentration on the reproduction of largemouth bass and bluegills. Prog. Fish-Cult. 26:99-106.
- Trama, F. B. 1954. The pH tolerance of the common bluegill, Lepomis macrochirus Raf. Not. Nat. (Phil.) 256:1-13.
- Trautman, M. B. 1957. The fishes of Ohio. Ohio State Univ. Press, Columbus. 683 pp.
- Ultsch, G. R. 1978. Oxygen consumption as a function of pH in three species of freshwater fishes. Copeia 1978:272-279.
- Weaver, R. O., and C. D. Ziebell. 1976. Ecology and early life history of largemouth bass and bluegill in Imperial Reservoir, Arizona. South-western Nat. 21:151-160.
- Werner, R. G. 1969. Ecology of limnetic bluegill (Lepomis macrochirus) fry in Crane Lake, Indiana. Am. Midl. Nat. 8:164-181.
- Whitmore, C. M., C. E. Warren, and P. Doudoroff. 1960. Avoidance reactions of salmonid and centrarchid fishes to low oxygen concentrations. Trans. Am. Fish. Soc. 89:17-26.

REPORT DOCUMENTATION PAGE	1. REPORT NO. FWS/OBS-82/10.8	2.	3. Recipient's Accession No.
4. Title and Subtitle Habitat Suitability Index Models: Bluegill			5. Report Date February 1982
7. Author(s) Robert J. Stuber, Glen Gebhart, and O. Eugene Maughan			6.
9. Performing Organization Name and Address Oklahoma Cooperative Fishery Research Unit Stillwater, OK 74074			8. Performing Organization Rept. No.
12. Sponsoring Organization Name and Address Western Energy and Land Use Team Office of Biological Services Fish and Wildlife Service U.S. Department of Interior Washington, D.C. 20240			10. Project/Task/Work Unit No.
15. Supplementary Notes			11. Contract(C) or Grant(G) No. (C) (G)
16. Abstract (Limit: 200 words) A literature review encompassing habitat and species characteristics of the bluegill (<i>Lepomis macrochirus</i>) is followed by a discussion of the relationship of habitat variables and life requisites of this species. These data are then incorporated into Habitat Suitability Index models for the bluegill. This is one in a series of publications describing habitat requirements of selected fish and wildlife species. Numerous literature sources have been consulted in an effort to consolidate scientific data on species habitat relationships. These data have subsequently been synthesized into Habitat Suitability Index (HSI) models. The models are based on suitability indices formulated for variables found to affect the life cycle and survival of the species. The models are designed to be modified to evaluate specific habitat alterations using the HSI model building techniques presented in the U.S. Fish and Wildlife Service's Habitat Evaluation Procedures.			13. Type of Report & Period Covered
17. Document Analysis a. Descriptors Habitat Suitability Fishes			14.
b. Identifiers/Open-Ended Terms Bluegill Lepomis macrochirus Habitat Habitat Suitability Index			
c. COSATI Field/Group			
18. Availability Statement Unlimited		19. Security Class (This Report) Unclassified	21. No. of Pages 26
		20. Security Class (This Page) Unclassified	22. Price



REGION 1

Regional Director
 U.S. Fish and Wildlife Service
 Lloyd Five Hundred Building, Suite 1692
 500 N.E. Multnomah Street
 Portland, Oregon 97232

REGION 2

Regional Director
 U.S. Fish and Wildlife Service
 P.O. Box 1306
 Albuquerque, New Mexico 87103

REGION 3

Regional Director
 U.S. Fish and Wildlife Service
 Federal Building, Fort Snelling
 Twin Cities, Minnesota 55111

REGION 4

Regional Director
 U.S. Fish and Wildlife Service
 Richard B. Russell Building
 75 Spring Street, S.W.
 Atlanta, Georgia 30303

REGION 5

Regional Director
 U.S. Fish and Wildlife Service
 One Gateway Center
 Newton Corner, Massachusetts 02158

REGION 6

Regional Director
 U.S. Fish and Wildlife Service
 P.O. Box 25486
 Denver Federal Center
 Denver, Colorado 80225

REGION 7

Regional Director
 U.S. Fish and Wildlife Service
 1011 E. Tudor Road
 Anchorage, Alaska 99503



DEPARTMENT OF THE INTERIOR U.S. FISH AND WILDLIFE SERVICE



As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally owned public lands and natural resources. This includes fostering the wisest use of our land and water resources, protecting our fish and wildlife, preserving the environmental and cultural values of our national parks and historical places, and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to assure that their development is in the best interests of all our people. The Department also has a major responsibility for American Indian reservation communities and for people who live in island territories under U.S. administration.